

Nohak[®]

Expectorant

COMPOSITION :

Nohak[®] Syrup: Each 5 ml syrup contains Guaifenesin BP 100 mg, Levomenthol BP 1.1 mg and Diphenhydramine HCL BP 14 mg.

PHARMACOLOGY :

Nohak[®] is a combined preparation of expectorant, decongestant and antihistamine. Guaifenesin is an expectorant. It helps to loosen mucus in chest and throat caused by common cold, infections, or allergies. Levomenthol helps to breathe more easily by relieving congestion. Diphenhydramine is an antihistamine that reduces the effects of natural chemical histamine in the body. It is used to treat sneezing, runny nose, watery eyes, hives, skin rash, itching & other cold or allergy symptoms.

INDICATION :

Nohak[®] is indicated for the symptomatic relief of upper-respiratory tract disorders accompanied by productive cough.

DOSAGE AND ADMINISTRATION :

12 years and above : 10 ml (2 teaspoonfuls) 4 times daily.

CONTRAINDICATION :

This combination is contraindicated in patients with a known hypersensitivity to Guaifenesin, Levomenthol or Diphenhydramine. If Monoamine Oxidase Inhibitors (MAOIs) have been taken in the last two weeks this combination should not be taken.

WARNING & PRECAUTION :

Patients suffering from liver, kidney & prostate problems should avoid this combination. Alcohol must be avoided. If any other medicines like minor Tranquilisers, Barbiturates, Codeine are taken, caution should be exercised. Patients taking this combination need to be careful while driving or operating machinery.

SIDE EFFECTS :

Side effects may include swelling of the face, lips and/or tongue, allergic reactions, numbness, drowsiness, dizziness, blurred vision, dry mouth/nose/throat, nausea or vomiting etc.

USE IN PREGNANCY AND LACTATION :

There are no specific data on use of this combination during pregnancy & lactation.

USE IN CHILDREN & ADOLESCENTS :

This preparation is not indicated for children under 12 years of age.

DRUG INTERACTION :

Taking certain Monoamine Oxidase Inhibitors (MAOIs) with this medication may cause a serious (possibly fatal) drug interaction. If any other medicines like minor Tranquilizers, Barbiturates and Codeine are taken, caution should be exercised.

OVERDOSE :

An overdose of this preparation may result in drowsiness, blurred vision, increased heart rate and confusion. Other symptoms may include dry mouth, nausea, vomiting, unsteadiness, high blood pressure and hallucinations.

STORAGE :

Protect from light, store in a cool and dry place below 30^o C temperature. Keep out of the reach of children.

PACKING :

Nohak[®] Syrup: Each bottle contains 100 ml syrup with a 10 ml measuring cup.



Manufactured by

Team Pharmaceuticals Ltd.

B 75-79, BSCIC, Rajshahi, Bangladesh